

$$\begin{array}{c}
 \begin{array}{c}
 \text{1. } \frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6} \\
 \text{2. } \frac{1}{4} \cdot \frac{1}{5} = \frac{1}{20} \\
 \text{3. } \frac{1}{6} \cdot \frac{1}{7} = \frac{1}{42} \\
 \text{4. } \frac{1}{8} \cdot \frac{1}{9} = \frac{1}{72} \\
 \text{5. } \frac{1}{10} \cdot \frac{1}{11} = \frac{1}{110} \\
 \text{6. } \frac{1}{12} \cdot \frac{1}{13} = \frac{1}{156} \\
 \text{7. } \frac{1}{14} \cdot \frac{1}{15} = \frac{1}{210} \\
 \text{8. } \frac{1}{16} \cdot \frac{1}{17} = \frac{1}{272} \\
 \text{9. } \frac{1}{18} \cdot \frac{1}{19} = \frac{1}{342} \\
 \text{10. } \frac{1}{20} \cdot \frac{1}{21} = \frac{1}{420}
 \end{array}
 \end{array}$$

Figure 1. The effect of the concentration of the *Agaricus bisporus* spores on the growth of *Agaricus bisporus* on the substrate. The concentration of the spores was 10⁵ spores/ml (a), 10⁶ spores/ml (b), 10⁷ spores/ml (c), 10⁸ spores/ml (d), 10⁹ spores/ml (e), 10¹⁰ spores/ml (f), 10¹¹ spores/ml (g), 10¹² spores/ml (h), 10¹³ spores/ml (i), 10¹⁴ spores/ml (j), 10¹⁵ spores/ml (k), 10¹⁶ spores/ml (l), 10¹⁷ spores/ml (m), 10¹⁸ spores/ml (n), 10¹⁹ spores/ml (o), 10²⁰ spores/ml (p), 10²¹ spores/ml (q), 10²² spores/ml (r), 10²³ spores/ml (s), 10²⁴ spores/ml (t), 10²⁵ spores/ml (u), 10²⁶ spores/ml (v), 10²⁷ spores/ml (w), 10²⁸ spores/ml (x), 10²⁹ spores/ml (y), 10³⁰ spores/ml (z).

10

© 2006 Blackwell Publishing Ltd, *Journal of Internal Medicine* 260: 101–108

30.1917
13
